

PROFILE FOR

St.Barnabas Surgery

St.Barnabas Terrace, Stoke, Plymouth

These profiles are designed to support clinical commissioning groups (CCGs), GPs and local authorities to ensure that they are providing and commissioning effective and appropriate healthcare services for their local population. The tool presents a range of practicelevel indicators drawn from the latest available data, including:

- local demography;
- Quality and Outcomes Framework (QOF) domains;
- patient satisfaction survey;
- cancer services

In addition to displaying individual practice profiles, the web tool allows you to view summary profiles for CCGs. Each practice can be compared with its CCG and with England, and also with the practices in the same deprivation deciles.

The profiles do not provide an exhaustive list of primary care indicators, but they do allow a consistent approach to comparing and benchmarking across England.


Note: QOF indicators are calculated

as percent of patients receiving an

intervention.

The profiles have been designed as a web tool and the full functionality and various chart types such as scatter plots and trend charts are only available via the web version. For more information consult the User guide and FAQs via the Supporting documents link, and for full metadata view the 'Definitions' on the website.

The development of this tool has been led by the Public Health England Fingertips team. For further information contact: ProfileFeedback@phe.gov.uk



http://fingertips.phe.org.uk/profile/general-practice

How to read the indicator spine charts

The light grey bar shows the range of values found in England. The dark grey sections mark out the range within which the middle half of the observed values lie (25th to 75th percentile).

The red line shows where the England average is. The position of the circle shows the practice value, a triangle the CCG value, in relation to this scale.

The corresponding numbers can be found in the cells next to the chart.

If significance has been calculated for the indicator, then it is determined by whether the practice value is significantly higher or lower than the England average usually using 99.8% confidence intervals.

- No significant difference from England average Significantly different from England average Significance not calculated **O** Practice
 - ▽ Clinical Commissioning Group

England Average England Highest England Lowest 75th Percentile 25th Percentile

| Practice Summary | | | | | | | | | | |
|--|-----------|----------------|--------------|--------------|--------------|-------------------------------|---------------|--|--|--|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. | | | |
| % aged 0 to 4 years | 2015 | 5.3% | 5.2% | 5.9% | 0.0% | $\overline{\mathbf{Q}}$ | 17.3% | | | |
| % aged 5 to 14 years | 2015 | 8.8% | 10.4% | 11.4% | 0.0% | | 30.3% | | | |
| % aged under 18 years | 2015 | 17.0% | 19.3% | 20.7% | 0.0% | | 53.5% | | | |
| % aged 65+ years | 2015 | 23.4% | 22.2% | 17.1% | 0.0% | | 92.5% | | | |
| % aged 75+ years | 2015 | 10.6% | 10.2% | 7.8% | 0.0% | \bigtriangledown | 79.6% | | | |
| % aged 85+ years | 2015 | 3.6% | 3.1% | 2.3% | 0.0% | \bigtriangledown | 48.2% | | | |
| Deprivation score (IMD 2015) | 2015 | 31.1 | 19.9 | 21.8 | 3.2 | \triangleleft O | 66.5 | | | |
| Deprivation score (IMD 2010) | 2012 | 30.0 | 19.8 | 21.5 | 2.9 | V O | 68.4 | | | |
| IDACI (Income Depr Children) | 2015 | 22.8% | 16.1% | 19.9% | 1.4% | $\nabla \bigcirc$ | 59.3% | | | |
| IDAOPI (Income Depr Older People) | 2015 | 21.8% | 13.6% | 16.2% | 3.9% | V O | 65.3% | | | |
| % who would recommend practice | 2014/15 | 86.5% | 85.6% | 77.5% | 15.2% | | 100% | | | |
| % satisfied with phone access | 2014/15 | 95.1% | 84.4% | 73.3% | 11.9% | | 100% | | | |
| % satisfied with opening hours | 2014/15 | 85.5% | 77.5% | 74.9% | 38.7% | \bigtriangledown | 100% | | | |
| % who saw/spoke to nurse or GP same or next day | 2014/15 | 49.7% | 49.3% | 48.3% | 6.5% | \Diamond | 98.3% | | | |
| % reporting good overall experience of making appointment | 2014/15 | 86.2% | 83.3% | 73.3% | 16.8% | | 100% | | | |
| % who know how to contact an out-of-hours GP service | 2014/15 | 69.0% | 66.1% | 56.4% | 11.9% | | 87.3% | | | |
| % with a long-standing health condition | 2014/15 | 55.9% | 56.5% | 54.0% | 11.9% | \bigtriangledown | 94.7% | | | |
| % with caring responsibility | 2014/15 | 15.2% | 19.2% | 18.2% | 0.5% | \bigcirc | 36.9% | | | |
| Working status - Paid work or full-time education | 2014/15 | 45.7% | 59.1% | 61.5% | 8.5% | $\bigcirc \forall$ | 100% | | | |
| Working status - Unemployed | 2014/15 | 4.3% | 4.0% | 5.4% | 0.4% | \bigcirc | 53.6% | | | |
| Total QOF points | 2014/15 | 95.3% | 95.1% | 94.8% | 28.2% | \bigtriangledown | 100% | | | |
| Life expectancy - MSOA based (Male) | 2008 - 12 | 75.2 | 79.6 | 78.9 | 70.0 | \bigcirc \bigtriangledown | 90.1 | | | |
| Life expectancy - MSOA based (Female) | 2008 - 12 | 79.9 | 83.4 | 82.8 | 75.9 | | 91.9 | | | |
| Nursing home patients | 2014/15 | 2.3% | 0.7% | 0.5% | 0.0% | $\checkmark \bigcirc$ | 14.9% | | | |

| | | С | ancer | | | | |
|--|---------|----------------|--------------|--------------|--------------|--------------------|---------------|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
| Cancer: QOF prevalence (all ages) | 2014/15 | 3.7% | 2.7% | 2.3% | 0.0% | | 6.2% |
| Exception rate for cancer indicator | 2014/15 | 30.8% | 17.1% | 15.4% | 0.0% | \bigtriangledown | 100% |
| New cancer cases (Crude incidence rate: new cases per 100,000 population) | 2012/13 | 639 | 631 | 508 | 0 | | 1593 |
| % reporting cancer in the last 5 years | 2014/15 | 4.5% | 3.8% | 3.3% | 0.3% | | 11.9% |
| CAN003: review within 6 mths of diagnosis | 2014/15 | 69.2% | 79.4% | 80.1% | 0.0% | \bigtriangledown | 100% |
| Exception rate for the cervical screening indicator | 2014/15 | 9.5% | 5.2% | 6.2% | 0.0% | \checkmark | 62.2% |
| CS002: Women, aged 25-64, with a record of cervical screening (last 5 yrs) (Female) | 2014/15 | 68.5% | 78.0% | 76.7% | 0.0% | | 100% |
| Females, 25-64, attending cervical screening within target period (3.5 or 5.5 year coverage, %) (Female) | 2014/15 | 73.2% | 76.9% | 73.5% | 0.0% | $\langle \nabla$ | 100% |
| Females, 50-70, screened for breast cancer in last 36 months (3 year coverage, %) (Female) | 2014/15 | 77.5% | 78.1% | 72.2% | 0.0% | | 94.1% |
| Females, 50-70, screened for breast cancer within 6 months of invitation (Uptake, %) (Female) | 2014/15 | 75.8% | 77.1% | 72.8% | 0.0% | \bigtriangledown | 100% |
| Persons, 60-69, screened for bowel cancer within 6 months of invitation (Uptake, %) | 2014/15 | 68.3% | 62.0% | 57.6% | 0.0% | | 100% |
| Persons, 60-69, screened for bowel cancer in last 30 months (2.5 year coverage, %) | 2014/15 | 65.7% | 62.7% | 57.9% | 10.1% | | 100% |
| Two-week wait referrals (Indirectly age-sex standardised referral ratio) | 2014/15 | 146 | 1 | 100 | 0 | \bigtriangledown | 614 |
| Number of new cancer cases treated (Detection rate: % of which resulted from a TWW referral) | 2014/15 | 58.8% | 53.3% | 48.4% | 0.0% | | 100% |
| Two-week wait referrals for suspected breast cancer (Number per 100,000 population) | 2014/15 | 761 | 595 | 482 | 0 | | 1645 |
| Two-week wait referrals for suspected lower GI cancers (Number per 100,000 population) | 2014/15 | 585 | 529 | 421 | 0 | | 2176 |
| Two-week wait referrals for suspected lung cancer (Number per 100,000 population) | 2014/15 | 117.0 | 110.0 | 100.5 | 0.0 | \diamond | 824.7 |
| Two-week wait referrals for suspected skin cancer (Number per 100,000 population) | 2014/15 | 1287 | 824 | 508 | 0 | | 3465 |
| In-patient or day-case colonoscopy procedures (Number per 100,000 population) | 2014/15 | 1170 | 806 | 695 | 0 | | 2105 |
| In-patient or day-case sigmoidoscopy procedures (Number per 100,000 population) | 2014/15 | 293 | 422 | 431 | 0 | \bigtriangledown | 1922 |
| In-patient or day-case upper GI endoscopy procedures (Number per 100,000 population) | 2014/15 | 878 | 1053 | 1142 | 28 | | 3576 |
| Number of emergency admissions with cancer (Number per 100,000 population) | 2014/15 | 936 | 549 | 539 | 6 | Ϋ́ | 2970 |
| Number of emergency presentations (Number per 100,000 population) | 2014/15 | 59 | 104 | 90 | 0 | \bigcirc | 721 |
| Number of other presentations (Number per 100,000 population) | 2014/15 | 761 | 491 | 360 | 0 | | 1186 |

CVD - Coronary heart disease

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|--|---------|----------------|--------------|--------------|--------------|--------------------|---------------|
| CHD: QOF prevalence (all ages) | 2014/15 | 6.1% | 3.9% | 3.2% | 0.0% | | 20.9% |
| Heart failure w LVD: QOF prevalence | 2014/15 | 0.4% | 0.2% | 0.2% | 0.0% | \checkmark | 2.2% |
| Exception rate for CHD indicators (2014/15 indicators) | 2014/15 | 6.2% | 9.4% | 8.4% | 0.0% | \Diamond | 66.7% |
| CHD002: Last BP reading in last 12mths is <=150/90 (den.incl.exc.) | 2014/15 | 89.4% | 87.3% | 88.4% | 0.0% | V | 100% |
| CHD003: Last total cholesterol is <=5mmol/l (den. incl. exc.) - retired | 2013/14 | 82.1% | 70.3% | 73.2% | 0.0% | VO | 100% |
| CHD005: Record that aspirin, APT or ACT is taken (den. incl. exc.) | 2014/15 | 96.2% | 92.0% | 91.7% | 0.0% | \bigtriangledown | 100% |
| CHD007: CHD patients immunised against flu (den.incl.exc.) | 2014/15 | 83.7% | 80.1% | 81.2% | 0.0% | \bigtriangledown | 100% |
| CHD006: History of MI: treated with ACE-I (den. incl. exc.) | 2014/15 | 77.8% | 64.0% | 69.1% | 0.0% | $\overline{\lor}$ | 100% |

CVD - Stroke and TIA

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|--|---------|----------------|--------------|--------------|--------------|-----------------------------|---------------|
| Stroke: QOF prevalence (all ages) | 2014/15 | 3.5% | 2.2% | 1.7% | 0.0% | | 20.3% |
| Exception rate for stroke indicators (2014/15 indicators) | 2014/15 | 2.2% | 10.4% | 9.7% | 0.0% | $\bigcirc \bigtriangledown$ | 66.7% |
| Exception rate for PAD indicators (2014/15 indicators) | 2014/15 | 2.2% | 6.6% | 5.8% | 0.0% | \bigcirc | 73.9% |
| STIA003: Last BP reading is =150/90 (den. incl. exc.) | 2014/15 | 91.5% | 83.6% | 84.3% | 0.0% | \bigtriangledown | 100% |
| STIA004: Total cholesterol recorded in last 15mths (den. incl. exc.) - retired | 2013/14 | 88.3% | 82.6% | 84.5% | 0.0% | \bigtriangledown | 100% |
| STIA005: Last measured total cholesterol <=5mmol/l (den. incl. exc.) - retired | 2013/14 | 81.5% | 64.4% | 68.4% | 0.0% | \triangleleft | 100% |
| STIA007: Record that an anti-platelet agent or an anti-coagulant is taken (den. incl. exc.) | 2014/15 | 100% | 92.9% | 91.7% | 0.0% | \checkmark | 100% |
| STIA008: New patients referred for further investigation (den.incl.exc.) | 2014/15 | 100% | 75.0% | 76.6% | 0.0% | \checkmark | 100% |
| STIA009: Influenza immunisation given 1 Aug-31 Mar (den.incl.exc.) | 2014/15 | 83.1% | 77.3% | 77.5% | 0.0% | \checkmark | 100% |
| PAD002: last blood pressure reading (last 12 mnths) <=150/90 mmHg (den.incl.exc.) | 2014/15 | 82.6% | 84.0% | 85.6% | 0.0% | V | 100% |
| PAD003: last total cholesterol measurement (last 12 mnths) <= 5.0 mmol/l (den.incl.exc.) - retired | 2013/14 | 84.6% | 66.1% | 69.4% | 0.0% | V O | 100% |
| PAD004: record of aspirin being taken (last 12 mnth) (den.incl.exc.) | 2014/15 | 100% | 88.1% | 86.4% | 0.0% | \bigtriangledown | 100% |

CVD - Heart failure and atrial fibrillation

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|---|---------|----------------|--------------|--------------|--------------|--------------------|---------------|
| Heart Failure: QOF prevalence (all ages) | 2014/15 | 0.9% | 0.8% | 0.7% | 0.0% | \bigtriangledown | 6.0% |
| Heart failure w LVD: QOF prevalence | 2014/15 | 0.4% | 0.2% | 0.2% | 0.0% | \checkmark | 2.2% |
| Exception rate for heart failure indicators | 2014/15 | 3.7% | 12.5% | 9.2% | 0.0% | | 78.0% |
| Atrial fibrillation: QOF prevalence | 2014/15 | 1.8% | 2.2% | 1.6% | 0.0% | \bigtriangledown | 14.1% |
| Estimated prevalence of Atrial Fibrillation | 2013/14 | 3.3% | 3.0% | 2.4% | 0.0% | \bigtriangledown | 16.7% |
| Estimated prevalence of Atrial Fibrillation (Male) | 2013/14 | 3.4% | 3.5% | | 0.1% | | 17.2% |
| Estimated prevalence of Atrial Fibrillation (Female) | 2013/14 | 3.2% | 2.5% | | 0.0% | | 16.4% |
| Estimated percentage of detected Atrial Fibrillation | 2013/14 | 65.5 | 68.9 | 65.2 | 0.0 | Ŷ | 131.9 |
| Exception rate for atrial fibrillation indicators (2014/15 indicators) | 2014/15 | 5.6% | 11.4% | 11.0% | 0.0% | \bigcirc | 100% |
| HF002: Diagnosis conf. by ECG/specialist assessm. (den. incl. exc.) | 2014/15 | 84.6% | 88.1% | 90.9% | 0.0% | V | 100% |
| HF003: Heart failure w LVD: treated with ACE-I or ARB (den. incl. exc.) | 2014/15 | 100% | 83.1% | 86.0% | 0.0% | $\forall igodom$ | 100% |
| HF004: Heart failure w LVD: treatment w ACE inh. or ARB, and beta-blocker (den.incl.exc.) | 2014/15 | 100% | 70.3% | 76.3% | 0.0% | \bigtriangledown | 100% |
| AF002: stroke risk assessed with CHADS2 (last 12 mnths) (den.incl.exc.) - retired | 2013/14 | 96.2% | 92.7% | 95.0% | 0.0% | ♥ | 100% |
| AF004: treated w anti-coag. therapy (CHADS2 >1) (den.incl.exc.) | 2014/15 | 93.8% | 77.0% | 74.3% | 0.0% | | 100% |
| AF005: treated w anti-coag./platelet therapy (if CHADS2 =1) (den.incl.exc.) | 2014/15 | 100% | 92.4% | 92.2% | 0.0% | \checkmark | 100% |

| | CVD | - Risk | factor | s for C | VD | | |
|--|---------|----------------|--------------|--------------|--------------|-----------------------------|---------------|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
| Hypertension: QOF prevalence (all ages) | 2014/15 | 15.4% | 15.2% | 13.8% | 0.1% | \bigtriangledown | 47.6% |
| Exception rate for hypertension indicator (2014/15) | 2014/15 | 1.1% | 5.0% | 3.8% | 0.0% | | 46.9% |
| Exception rate for the BP indicator | 2014/15 | 1.4% | 0.6% | 0.4% | 0.0% | \Diamond | 49.9% |
| Obesity: QOF prevalence (16+) | 2014/15 | 1.9% | 10.1% | 9.0% | 0.0% | | 29.4% |
| Estimated smoking prevalence (QOF) | 2014/15 | 22.3 | 17.3 | 18.4 | 0.0 | | 83.3 |
| Exception rate for smoking indicators (2014/15 indicators) | 2014/15 | 2.0% | 1.3% | 1.0% | 0.0% | \Diamond | 65.0% |
| GP patient survey: smoking prevalence | 2014/15 | 15.3% | 15.6% | 16.4% | 1.3% | \bigcirc | 85.8% |
| GP patient survey: ex-smoking prevalence | 2014/15 | 36.4% | 30.6% | 27.4% | 0.5% | \bigtriangledown | 51.3% |
| SMOK001: record of smoking status in last 24 months (15+ y), den. incl. exc retired | 2013/14 | 85.5% | 83.2% | 85.6% | 2.1% | $\overline{\Diamond}$ | 100% |
| SMOK002: status recorded in last 12 mths (certain conditions), den.incl.exc. | 2014/15 | 92.3% | 93.5% | 93.2% | 0.0% | \bigcirc | 100% |
| SMOK004: record of offer of support and treatment (15+, last 24 mnths), den. incl. exc. | 2014/15 | 86.8% | 84.9% | 85.8% | 0.0% | Ô | 100% |
| SMOK005: cessation support and treatment offered (certain conditions), den. incl. exc. | 2014/15 | 98.0% | 93.3% | 94.1% | 12.6% | Ż | 100% |
| HYP006: Blood pressure <= 150/90 mmHg in people with hypertension | 2014/15 | 87.5% | 80.2% | 80.4% | 0.0% | \checkmark | 100% |
| HYP003: Last (9mnths) blood pressure <= 140/90 mmHg (age <80) (den.incl.exc.) - retired | 2013/14 | 75.5% | 64.1% | 70.4% | 0.0% | $\nabla \bigcirc$ | 98.9% |
| HYP004: Phys. act. assessment (last 12 mnths), patients with hypertension aged 16-74 (den.incl.exc.) - retired | 2013/14 | 76.5% | 62.5% | 76.0% | 0.0% | $\bigtriangledown \bigcirc$ | 100% |
| HYP005: Patients w. hypertension who had a phys. act. assessment and where found to be inactive who had also a brief intervention (both in last 12mnths, aged 16-74) (den.incl.exc.) - retired | 2013/14 | 69.7% | 78.4% | 86.3% | 0.0% | | 100% |
| BP002: Patients, aged 45+, who have a record of blood pressure (last 5yrs) | 2014/15 | 91.4% | 91.1% | 90.6% | 0.0% | \heartsuit | 100% |
| CVD-PP001: new hypertension patients, age 30-74, with CV risk assessment >=20% treated w. statins (den.incl.exc.) | 2014/15 | - | 65.5% | 67.7% | 0.0% | | 100% |
| CVD-PP002: Lifestyle advice for patients with hypertension (den. incl. exc.) - retired | 2013/14 | 68.2% | 66.4% | 79.0% | 0.0% | | 100% |

| | | Chil | d heal | th | | | |
|---|--------------------|----------------|--------------|--------------|--------------|--------------------|---------------|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
| % aged 0 to 4 years | 2015 | 5.3% | 5.2% | 5.9% | 0.0% | \bigcirc | 17.3% |
| % aged 5 to 14 years | 2015 | 8.8% | 10.4% | 11.4% | 0.0% | | 30.3% |
| % aged under 18 years | 2015 | 17.0% | 19.3% | 20.7% | 0.0% | | 53.5% |
| IDACI (Income Depr Children) | 2015 | 22.8% | 16.1% | 19.9% | 1.4% | ∇O | 59.3% |
| A&E attendances (0-4) | 2011/12 - 13/14 | 204 | 471 | 530 | 22 | O.↓ | 2705 |
| A&E attendances (5-17) | 2011/12 - 13/14 | 211 | 301 | 307 | 5 | \bigtriangledown | 1145 |
| A&E attendances (<18) | 2011/12 - 13/14 | 209 | 347 | 373 | 11 | O.↓ | 1504 |
| Elective hospital admissions for all causes (<18) | 2011/12 - 13/14 | 30.5 | 47.7 | 49.0 | 1.6 | \bigtriangledown | 249.1 |
| Emergency hospital admissions for all causes (<18) | 2011/12 - 13/14 | 63.4 | 65.8 | 69.5 | 2.5 | Ø | 270.0 |
| Emergency respiratory admissions (<18) | 2009/10 - 13/14 | - | 3.8 | 3.8 | 0.4 | \bigtriangledown | 14.5 |
| Emergency gastroenteritis admissions (0-4) | 2009/10 - 13/14 | 20.8 | 11.4 | 11.3 | 0.8 | ΫO | 72.1 |
| Emergency admissions for asthma, diabetes or epilepsy (<18) | 2009/10 - 13/14 | - | 3.0 | 3.4 | 0.4 | Δ | 54.7 |
| Admissions due to injury (<18) | 2011/12 - 13/14 | - | 10.6 | 11.1 | 0.7 | $\dot{\nabla}$ | 52.8 |
| Outpatient first attendances (<18) | 2011/12 - 13/14 | 263 | 282 | 246 | 6 | \bigtriangledown | 1144 |
| Ratio of first to follow-up outpatient attendances (<18) | 2011/12 - 13/14 | 0.38 | 0.46 | 0.51 | 0.20 | \bigtriangledown | 2.14 |
| DNA rate for outpatient appointments (<18) | 2011/12 - 13/14 | 7.5% | 10.7% | 11.5% | 0.6% | $\bigcirc \forall$ | 32.3% |

Chronic Kidney Disease

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|---|---------|----------------|--------------|--------------|--------------|-------------------------|---------------|
| CKD: QOF prevalence (18+) | 2014/15 | 8.2% | 4.9% | 4.1% | 0.0% | \bigtriangledown | 23.8% |
| Exception rate for CKD indicators | 2014/15 | 3.1% | 9.5% | 7.5% | 0.0% | | 60.0% |
| CKD002: Last BP reading measured in last 12mths is <=140/85 (den. incl. exc.) | 2014/15 | 80.3% | 71.1% | 74.4% | 0.0% | $\overline{\mathbf{v}}$ | 100% |
| CKD003: Hypertension treated with ACE inhibitor/ARB (den. incl. exc.) | 2014/15 | 91.3% | 76.0% | 76.4% | 0.0% | \ | 100% |
| CKD004: Urine albumin:creatinine ratio test last 12 mths (den. incl. exc.) | 2014/15 | 76.9% | 75.8% | 75.4% | 0.0% | \bigtriangledown | 100% |

| Diabetes | | | | | | | | | |
|---|---------|----------------|--------------|--------------|--------------|--------------------|---------------|--|--|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. | | |
| Diabetes: QOF prevalence (17+) | 2014/15 | 8.0% | 6.5% | 6.4% | 0.3% | \bigtriangledown | 20.4% | | |
| Exception rate for diabetes indicators (2014/15 indicators) | 2014/15 | 10.1% | 12.6% | 10.8% | 0.0% | \bigcirc | 49.0% | | |
| Hypertension: QOF prevalence (all ages) | 2014/15 | 15.4% | 15.2% | 13.8% | 0.1% | \bigtriangledown | 47.6% | | |
| Exception rate for hypertension indicator (2014/15) | 2014/15 | 1.1% | 5.0% | 3.8% | 0.0% | \bigcirc | 46.9% | | |
| Exception rate for smoking indicators (2014/15 indicators) | 2014/15 | 2.0% | 1.3% | 1.0% | 0.0% | \diamond | 65.0% | | |
| Obesity: QOF prevalence (16+) | 2014/15 | 1.9% | 10.1% | 9.0% | 0.0% | | 29.4% | | |
| DM007: Last HbA1c is <=59mmol/mol in last 12mths (den. incl. exc.) | 2014/15 | 67.2% | 60.3% | 60.4% | 0.0% | V. | 100% | | |
| DM008: Last HbA1c is <=64mmol/mol in last 12mths (den. incl. exc.) | 2014/15 | 70.7% | 68.7% | 68.4% | 0.0% | \bigtriangledown | 100% | | |
| DM009: Last HbA1c is <=75mmol/mol in last 12mths (den. incl. exc.) | 2014/15 | 77.6% | 80.4% | 79.3% | 0.0% | \bigtriangledown | 100% | | |
| DM002: Last BP is <=150/90 (den. incl. exc.) | 2014/15 | 92.2% | 85.4% | 86.6% | 0.0% | \checkmark | 100% | | |
| DM003: Last BP is <=140/80 (den. incl. exc.) | 2014/15 | 85.3% | 67.2% | 71.2% | 0.0% | \bigtriangledown | 100% | | |
| DM004: Measured total cholesterol (last 12mths) <=5mmol/l(den.incl.exc.) | 2014/15 | 82.8% | 70.2% | 70.8% | 0.0% | \bigtriangledown | 100% | | |
| DM005: Record of micro-albuminuria test last 12mths (den. incl. exc.) - retired | 2013/14 | 78.0% | 76.7% | 80.6% | 1.1% | \Diamond | 100% | | |
| DM006: Proteinuria/micro-album. treated w inhibitors (den. incl. exc.) | 2014/15 | 92.9% | 79.2% | 81.2% | 0.0% | $\forall \bigcirc$ | 100% | | |
| DM011: record of retinal screening (den.incl.exc.) - retired | 2013/14 | 69.9% | 77.4% | 82.6% | 0.0% | | 100% | | |
| DM012: Patients with diabetes who had a foot examination and risk classification (den.incl.exc.) | 2014/15 | 75.0% | 81.7% | 81.5% | 0.0% | ⇔ | 100% | | |
| DM013: Patients with diabetes who had a dietary review (last 12mths) (den.incl.exc.) - retired | 2013/14 | 57.7% | 75.9% | 82.2% | 0.0% | | 100% | | |
| DM014: Newly diagnosed patients w. diabetes referred to education programme within 9 mths (den.incl.exc.) | 2014/15 | 16.7% | 49.6% | 66.3% | 0.0% | | 100% | | |
| DM015: Male patients w. diabetes asked about erectile disfunction (last 12 mths) (den.incl.exc.) - retired (Male) | 2013/14 | 77.3% | 69.8% | 81.6% | 0.0% | | 100% | | |
| DM016: Male patients w. diabetes w. erectile disfunction who got advice (den.incl.exc.) - retired (Male) | 2013/14 | 50.0% | 77.8% | 87.9% | 0.0% | | 100% | | |
| DM018: Flu vaccination coverage (den.incl.exc.) | 2014/15 | 73.3% | 76.4% | 77.6% | 37.0% | | 100% | | |

| | | Ment | al Hea | lth | | | |
|---|---------|----------------|--------------|--------------|--------------|-------------------------|---------------|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
| Mental Health: QOF prevalence (all ages) | 2014/15 | 1.99% | 0.85% | 0.88% | 0.06% | \checkmark | 15.57% |
| Exception rate for MH indicators (2014/15 indicators) | 2014/15 | 8.3% | 11.8% | 11.1% | 0.0% | \bigtriangledown | 76.7% |
| Dementia: QOF prevalence (all ages) | 2014/15 | 2.0% | 0.9% | 0.7% | 0.0% | \Diamond | 59.2% |
| % reporting Alzheimer's disease or dementia | 2014/15 | 2.1% | 0.9% | 1.0% | 0.3% | \bigtriangledown | 25.0% |
| Exception rate for dementia indicators | 2014/15 | 2.6% | 9.8% | 8.3% | 0.0% | \bigcirc | 100% |
| Depression: QOF incidence (18+) - new diagnosis | 2014/15 | 2.0 | 1.3 | 1.2 | 0.0 | \bigtriangledown | 18.2 |
| Depression: QOF prevalence (18+) | 2014/15 | 14.6% | 8.1% | 7.3% | 0.0% | \forall \bigcirc | 33.0% |
| Exception rate for depression indicator (2014/15 indicators) | 2014/15 | 37.9% | 25.5% | 24.5% | 0.0% | \Diamond | 100% |
| % reporting a long-term mental health problem | 2014/15 | 9.4% | 4.8% | 5.1% | 0.4% | \checkmark | 60.1% |
| MH002: comprehensive care plan (den. incl. exc.) | 2014/15 | 85.7% | 74.7% | 77.2% | 0.0% | ∀ ⊖ | 100% |
| MH007: record of alcohol consumption for patients on the MH register (last 12 mnths), den. incl. exc. | 2014/15 | 67.9% | 77.1% | 80.3% | 0.0% | | 100% |
| MH006: record of BMI for patients on the MH register in preceding 12 months (den.incl.exc.) - retired | 2013/14 | 87.1% | 75.3% | 78.8% | 0.0% | Ч <mark>О</mark> | 100% |
| MH003: record of blood pressure check in preceding 12 months for patients on the MH register (den.incl.exc.) | 2014/15 | 71.4% | 79.7% | 81.5% | 0.0% | | 100% |
| MH008: Female patients (25-64 yrs) on the MH register who had cervical screening test in preceding 5 years (den. incl. exc.) | 2014/15 | 50.0% | 71.5% | 71.6% | 0.0% | \bigcirc \Diamond | 100% |
| MH009: Patients on lithium therapy with record of serum creatinine and TSH in the preceding 9 months (den.incl.exc.) | 2014/15 | 100% | 93.9% | 93.4% | 0.0% | \checkmark | 100% |
| MH010: Patients on lithium therapy with levels in therapeutic range in preceding 4 months (den.incl.exc.) | 2014/15 | 66.7% | 84.0% | 82.6% | 0.0% | \bigcirc \checkmark | 100% |
| MH004: Patients on MH register with cholesterol check in preceding 12 months (den.incl.exc.) - retired | 2013/14 | 80.0% | 61.4% | 68.0% | 0.0% | $\nabla \bigcirc$ | 100% |
| MH005: Patients on the MH register with blood glucose or HbA1c check in preceding 12 months (den.incl.exc.) - retired | 2013/14 | 76.2% | 71.0% | 74.9% | 0.0% | \checkmark | 100% |
| DEP001: Newly diagnosed patients w. depression who had a bio-psychosocial assessment on diagnosis (current FY, aged 18+) (den.incl.exc.) - retired | 2013/14 | 80.0% | 68.3% | 75.8% | 0.0% | ∇ <mark></mark> | 100% |
| DEP003: Newly diagnosed patients with depression who had a review 10-56 days after diagnosis (den.incl.exc.) | 2014/15 | 58.6% | 63.4% | 63.8% | 0.0% | Ċ | 100% |
| DEM002: Dementia care has been reviewed last 12mths (den.incl.exc.) | 2014/15 | 73.5% | 77.9% | 77.0% | 0.0% | ○ | 100% |
| DEM003: Blood tests recorded (den.incl.exc.) | 2014/15 | 100% | 73.1% | 74.7% | 0.0% | V O | 100% |

Musculoskeletal Conditions

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|---|---------|----------------|--------------|--------------|--------------|--------------------|---------------|
| Osteoporosis: QOF prevalence (50+) | 2014/15 | 0.5 | 0.2 | 0.2 | 0.0 | \checkmark | 9.6 |
| Exception rate for OST indicators | 2014/15 | 0.0% | 13.0% | 12.5% | 0.0% | $\bigcirc \forall$ | 100% |
| Rheumatoid Arthritis: QOF prevalence (16+) | 2014/15 | 0.55 | 0.82 | 0.73 | 0.00 | \triangleleft | 4.60 |
| Exception rate for RA indicators (2014/15 indicators) | 2014/15 | 12.5% | 8.4% | 7.4% | 0.0% | \checkmark | 82.8% |
| % reporting a long-term back problem | 2014/15 | 11.7% | 9.6% | 9.9% | 0.0% | \checkmark | 38.4% |
| % reporting arthritis or long-term joint problem | 2014/15 | 16.4% | 13.4% | 12.8% | 0.4% | ∇ | 34.9% |
| OST002: currently treated with a bone- sparing agent (50-74 yrs)(den.incl.exc.) | 2014/15 | 100% | 82.7% | 82.8% | 0.0% | \Diamond | 100% |
| OST005: Patients (75+ yrs) with a fragility fracture treated with bone-sparing agent (den.incl.exc.) | 2014/15 | 100% | 78.4% | 79.3% | 0.0% | Ý 🕒 | 100% |
| RA002: Patients with rheumatoid arthritis who had a face-to-face review (last 12 mnths) (den.incl.exc.) | 2014/15 | 37.5% | 84.7% | 84.3% | 0.0% | | 100% |
| RA003: Patients w. rheumatoid arthritis, aged 30-84, who had an RA adjusted CV risk assessment (last 12 mnths) (den.incl.exc.) - retired | 2013/14 | 100% | 75.0% | 86.8% | 0.0% | ∇ | 100% |
| RA004: Patients w. rheumatoid arthritis, aged 50-90, who had an RA adjusted fracture risk assessment (last 24 mnths) (den.incl.exc.) - retired | 2013/14 | 100% | 68.5% | 82.1% | 0.0% | | 100% |

| Respiratory Disease | | | | | | | | | |
|---|---------|----------------|--------------|--------------|--------------|--------------------|---------------|--|--|
| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. | | |
| COPD: QOF prevalence (all ages) | 2014/15 | 2.4% | 2.0% | 1.8% | 0.0% | \bigtriangledown | 9.0% | | |
| Exception rate for COPD indicators | 2014/15 | 13.1% | 12.6% | 12.3% | 0.0% | \bigtriangledown | 57.8% | | |
| Asthma: QOF prevalence (all ages) | 2014/15 | 8.0% | 6.7% | 6.0% | 0.7% | | 12.5% | | |
| Exception rate for asthma indicators | 2014/15 | 4.2% | 9.4% | 6.8% | 0.0% | | 63.7% | | |
| Estimated smoking prevalence (QOF) | 2014/15 | 22.3 | 17.3 | 18.4 | 0.0 | \checkmark | 83.3 | | |
| Exception rate for smoking indicators (2014/15 indicators) | 2014/15 | 2.0% | 1.3% | 1.0% | 0.0% | \Diamond | 65.0% | | |
| GP patient survey: smoking prevalence | 2014/15 | 15.3% | 15.6% | 16.4% | 1.3% | \bigcirc | 85.8% | | |
| GP patient survey: ex-smoking prevalence | 2014/15 | 36.4% | 30.6% | 27.4% | 0.5% | | 51.3% | | |
| COPD007: Influenza immunisation given 1 Aug - 31 Mar (den. incl. exc.) | 2014/15 | 85.4% | 80.8% | 81.5% | 0.0% | \bigtriangledown | 100% | | |
| COPD004: Record of FEV1 in last 12mths (den. incl. exc.) | 2014/15 | 78.0% | 75.8% | 73.2% | 0.0% | \bigtriangledown | 100% | | |
| COPD003: assessed using MRC dyspnoea score last 12mths (den. incl. exc.) | 2014/15 | 75.6% | 79.5% | 79.9% | 0.0% | ◯ | 100% | | |
| COPD002: Diagnosis conf. by spirometry (den. incl. exc.) | 2014/15 | 80.0% | 79.5% | 81.1% | 0.0% | \bigotimes | 100% | | |
| AST002: with measures of variability/reversibility (8+), den. incl. exc. | 2014/15 | 88.9% | 83.6% | 84.3% | 0.0% | \checkmark | 100% | | |
| AST003: review in the last 12 months (incl. an assessment using the 3 RCP questions), den. incl. exc. | 2014/15 | 70.6% | 67.8% | 69.7% | 0.0% | V | 100% | | |
| AST004: smoking recorded in last 12 mths (14-19y w asthma), den. incl. exc. | 2014/15 | 80.0% | 81.2% | 83.9% | 0.0% | V | 100% | | |
| SMOK002: status recorded in last 12 mths (certain conditions), den.incl.exc. | 2014/15 | 92.3% | 93.5% | 93.2% | 0.0% | V | 100% | | |
| SMOK005: cessation support and treatment offered (certain conditions), den. incl. exc. | 2014/15 | 98.0% | 93.3% | 94.1% | 12.6% | \checkmark | 100% | | |
| SMOK001: record of smoking status in last 24 months (15+ y), den. incl. exc retired | 2013/14 | 85.5% | 83.2% | 85.6% | 2.1% | V. | 100% | | |
| SMOK004: record of offer of support and treatment (15+, last 24 mnths), den. incl. exc. | 2014/15 | 86.8% | 84.9% | 85.8% | 0.0% | V | 100% | | |
| COPD005: Patients w. MRC dyspnoea score >=3 w.oxygen saturation value (last 12mths) (den.incl.exc.) | 2014/15 | 82.4% | 94.6% | 94.5% | 0.0% | ○ 文 | 100% | | |

AMR Local indicators

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|---|---------|----------------|--------------|--------------|--------------|--------------------|---------------|
| Total number of prescribed antibiotic items per 1000 registered patients by quarter | 2016 Q1 | 281.2 | 161.1 | 160.2 | 0.1 | \checkmark | 2119.7 |
| Total number of prescribed antibiotic items per STAR-PU by quarter | 2016 Q1 | 0.47 | 0.27 | 0.30 | 0.00 | \checkmark | 4.06 |
| Percentage of broad spectrum prescribed antibiotic items (cephalosporin, quinolone | 2016 Q1 | 15.04% | 10.96% | 9.87% | 0.00% | \bigtriangledown | 100.00% |

and co-amoxiclav class) by quarter

Other Conditions

| Indicator | Period | Prac. Value | CCG Value | Eng. Ave. | Eng. Low. | England Range | Eng. High. |
|---|---------|----------------|--------------|--------------|--------------|------------------------------|---------------|
| Epilepsy: QOF prevalence (18+) | 2014/15 | 1.7% | 0.9% | 0.8% | 0.1% | \bigtriangledown | 5.6% |
| EP002: Seizure free for last 12mths (den.incl.exc.) - retired | 2013/14 | 59.1% | 56.1% | 61.2% | 0.0% | $\overline{\mathbf{\nabla}}$ | 100% |
| EP003: Contraception, conception and pregnancy (den.incl.exc.) - retired (Female) | 2013/14 | 33.3% | 39.6% | 56.9% | 0.0% | | 100% |
| Learning disability: QOF prevalence | 2014/15 | 2.2% | 0.5% | 0.4% | 0.0% | \bigtriangledown | 4.3% |
| LD002: Down's Syndrome with blood TSH record (den.incl.exc.) - retired | 2013/14 | 40.0% | 48.2% | 54.7% | 0.0% | | 100% |
| % reporting learning difficulty | 2014/15 | 3.2% | 2.9% | 2.6% | 0.3% | \Diamond | 30.8% |
| Hypothyroidism: QOF prevalence (all ages) - retired | 2013/14 | 5.4% | 3.8% | 3.3% | 0.0% | | 10.8% |
| THY002: function test recorded last 12mths (den.incl.exc.) - retired | 2013/14 | 94.7% | 92.9% | 93.6% | 0.0% | Ø | 100% |
| Palliative/supportive care: QOF prevalence (all ages) | 2014/15 | 0.1% | 0.3% | 0.3% | 0.0% | \Diamond | 35.3% |
| % reporting blindness or severe visual impairment | 2014/15 | 2.9% | 1.4% | 1.4% | 0.3% | | 17.2% |
| % reporting deafness or severe hearing impairment | 2014/15 | 2.8% | 4.9% | 4.0% | 0.2% | | 17.2% |